

Effective Field Theory of Gravity: Leading Quantum Gravitational Corrections to Newtons and Coulombs Law

SVEN FALLER*

*Theoretische Physik 1, Fachbereich Physik, Universität Siegen
D-57068 Siegen, Germany*

In this paper we consider general relativity and its combination with scalar quantum electrodynamics (QED) as an effective quantum field theory at energies well below the Planck scale. This enables us to compute the one-loop quantum corrections to the Newton and Coulomb potential induced by the combination of graviton and photon fluctuations. We derive the relevant Feynman rules and compute the nonanalytical contributions to the one-loop scattering matrix for charged scalars in the nonrelativistic limit. In particular, we derive the post-Newtonian corrections of order $Gm/c^2 r$ from general relativity and the genuine quantum corrections of order $G\hbar/c^3 r^2$.

I. INTRODUCTION

At the classical level the interaction between two charged massive particles is described by Newton's and Coulomb's law, in terms of the nonrelativistic potential

$$V(r) = -G \frac{m_1 m_2}{r} + \frac{1}{4\pi} \frac{e_1 e_2}{r} . \quad (1)$$

This potential will be modified by relativistic and quantum corrections. Radiative corrections to the Coulomb potential can be systematically calculated within QED. However, it is well known that general relativity, including scalar [1, 2, 3], fermion or photon fields [4, 5, 6] is not renormalizable. Still, on smooth-enough background space-time the gravitational field can be quantized consistently [1, 2], and as suggested by Donoghue [7, 8], one can consider general relativity as an effective quantum field theory in the low-energy limit where renormalizability is no longer an issue.

Within this framework, the one-loop corrections to quantum gravity coupled to a scalar Klein-Gordon field have already been discussed by independent groups, e.g. Donoghue [7, 8, 9], Hamber and Liu [10], Akhundov et al. [11, 12], and Bjerrum-Bohr et al. [13, 14, 15], with differing results. The generalization to scalar QED has been first discussed by Bjerrum-Bohr [14]. Butt [16] has also treated the case of charged fermions.

In this paper, we are going to reconsider the results for the case of scalar QED coupled to gravity (SQED), in order to perform an independent check of the above results and to resolve the discrepancies between [7, 8, 9], [10], [11, 12] and [13, 14, 15]. To this end, we discuss in Sec. II how to quantize general relativity/SQED with the background field method. In the next section we review how ultraviolet divergences from one-loop diagrams

can be absorbed into higher-dimensional operators in the effective theory. The definition of the scattering matrix and of the potential for (charged or neutral) scalar particles is given in Sec. IV. In Sec. V we perform the calculations of the relevant Feynman diagrams and construct the scattering potentials at one loop. We compare our results with [7, 8, 11, 12, 13] and [14], before we conclude with a short summary. The Feynman rules, one-loop integrals, and details of the Fourier transformations are presented in the Appendices A, B and C.

II. QUANTIZATION OF GENERAL RELATIVITY AND SCALAR QED

The Einstein-Hilbert action for the vacuum is given by

$$\mathcal{S}_{\text{vac}} = \int d^4x \sqrt{-g} \frac{2}{\kappa^2} \mathcal{R} , \quad (2)$$

where $\kappa^2 = 32\pi G$ is the gravitational coupling, $g_{\mu\nu}$ is the metric tensor with its determinant $g = \det g_{\mu\nu}$, and $\mathcal{R} = g^{\mu\nu} \mathcal{R}_{\mu\nu}$ is the curvature scalar. In the following we use the same convention as Refs. [7, 8, 9, 11, 12, 13, 14, 15, 16], with the Minkowski metric $\eta_{\mu\nu} = \text{diag}(+1, -1, -1, -1)$ and $\hbar = c = 1$.

Additional fields can be included by adding a covariant term $\sqrt{-g} \mathcal{L}_m$ to the action Eq. (2),

$$\mathcal{S} = \int d^4x \sqrt{-g} \left[\frac{2}{\kappa^2} \mathcal{R} + \mathcal{L}_m \right] . \quad (3)$$

For instance, massive scalar fields are described by the scalar Lagrangian

$$\mathcal{L}_m = \frac{1}{2} (g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi^* - m^2 |\phi|^2) . \quad (4)$$

*faller@hep.physik.uni-siegen.de

A. Background Field Method

Following Veltman and 't Hooft [1, 2] we can expand the gravitational field $g_{\mu\nu}$ into the classical background field $\bar{g}_{\mu\nu}$ and a small quantum fluctuation $\kappa h_{\mu\nu}$:

$$\begin{aligned} g_{\mu\nu} &= \bar{g}_{\mu\nu} + \kappa h_{\mu\nu} , \\ g^{\mu\nu} &= \bar{g}^{\mu\nu} - \kappa h^{\mu\nu} + \kappa^2 h^\mu_\lambda h^{\lambda\nu} . \end{aligned} \quad (5)$$

Expanding the curvature to second order in the quantum field h , using

$$\sqrt{-g} = \sqrt{-\bar{g}} \left\{ 1 + \frac{\kappa}{2} h^\alpha_\alpha - \frac{\kappa^2}{4} h^\alpha_\beta h^\beta_\alpha + \frac{\kappa^2}{8} (h^\alpha_\alpha)^2 + \dots \right\} , \quad (6)$$

we obtain for the gravitational part of the Lagrangian

$$\mathcal{L}_{\text{grav}} = \sqrt{-\bar{g}} \left[\frac{2}{\kappa^2} \bar{\mathcal{R}} + \mathcal{L}_{\text{grav}}^{(1)} + \mathcal{L}_{\text{grav}}^{(2)} + \dots \right] \quad (7)$$

with

$$\begin{aligned} \mathcal{L}_{\text{grav}}^{(1)} &= \frac{1}{\kappa} h_{\mu\nu} [\bar{g}^{\mu\nu} \bar{\mathcal{R}} - 2 \bar{\mathcal{R}}^{\mu\nu}] , \\ \mathcal{L}_{\text{grav}}^{(2)} &= \frac{1}{2} \bar{\mathcal{D}}_\alpha h_{\mu\nu} \bar{\mathcal{D}}^\alpha h^{\mu\nu} - \frac{1}{2} \bar{\mathcal{D}}_\alpha h \bar{\mathcal{D}}^\alpha h \\ &\quad + \bar{\mathcal{D}}_\alpha h \bar{\mathcal{D}}_\beta h^{\alpha\beta} - \bar{\mathcal{D}}_\alpha h_{\mu\beta} \bar{\mathcal{D}}^\beta h^{\mu\alpha} \\ &\quad + \bar{\mathcal{R}} \left(\frac{1}{4} h^2 - \frac{1}{2} h_{\mu\nu} h^{\mu\nu} \right) \\ &\quad + \bar{\mathcal{R}}^{\mu\nu} (2 h^\lambda_\mu h_{\nu\lambda} - h h_{\mu\nu}) . \end{aligned} \quad (8)$$

Here the curvature scalar $\bar{\mathcal{R}}$ and the covariant derivative $\bar{\mathcal{D}}_\alpha$ are evaluated from the background metric $\bar{g}_{\mu\nu}$.

In order to quantize gravity we have to fix the gauge and introduce a Faddeev-Popov ghost field. From [1] we take the gauge fixing Lagrangian

$$\mathcal{L}_{\text{gf}} = \sqrt{-\bar{g}} \left\{ \left(\bar{\mathcal{D}}^\nu h_{\mu\nu} - \frac{1}{2} \bar{\mathcal{D}}_\mu h \right) \left(\bar{\mathcal{D}}_\lambda h^{\mu\lambda} - \frac{1}{2} \bar{\mathcal{D}}^\mu h \right) \right\} , \quad (9)$$

and the ghost Lagrangian

$$\mathcal{L}_{\text{ghost}} = \sqrt{-\bar{g}} \eta^{*\mu} [\bar{\mathcal{D}}_\lambda \bar{\mathcal{D}}^\lambda g_{\mu\nu} - \bar{\mathcal{R}}_{\mu\nu}] \eta^\nu , \quad (10)$$

with η^ν being the complex Faddeev-Popov ghost field. From this Lagrangian one derives the graviton propagator as quoted in Appendix A.

B. Combining Scalar QED and General Relativity

The generally covariant Lagrangian for scalar QED is [14]

$$\begin{aligned} \mathcal{L}_{\text{SQED}} &= \sqrt{-g} \left[-\frac{1}{4} (g^{\alpha\mu} g^{\beta\nu} \mathcal{F}_{\alpha\nu} \mathcal{F}_{\mu\beta}) \right. \\ &\quad \left. + g^{\mu\nu} \mathcal{D}_\mu \phi \mathcal{D}_\nu \phi^* - m^2 |\phi|^2 \right] , \end{aligned} \quad (11)$$

with the photon field strength $\mathcal{F}_{\mu\nu} = \mathcal{D}_\mu \mathcal{A}_\nu - \mathcal{D}_\nu \mathcal{A}_\mu$, and the QED covariant derivative $\mathcal{D}_\mu = \partial_\mu - ie_q \mathcal{A}_\mu(x)$, and e_q being the charge of the scalar particle. In order to get the relevant Feynman rules at one-loop order, we have to expand the Lagrangian $\mathcal{L}_{\text{SQED}}$ to first order in the quantum field h . Using Eqs. (5), (7) and working with the Lorenz gauge for the photon field,

$$\mathcal{L}^{\text{C}} = -\frac{1}{2} (\partial_\mu \mathcal{A}^\mu)^2 , \quad (12)$$

we get for the photon part

$$\begin{aligned} \mathcal{L}_{\text{photon}}^{(1)} &= \sqrt{-\bar{g}} \left\{ -\frac{\kappa h}{4} (\partial_\mu \mathcal{A}_\nu \partial^\mu \mathcal{A}^\nu - \partial_\mu \mathcal{A}_\nu \partial^\nu \mathcal{A}^\mu) \right. \\ &\quad + \frac{\kappa}{2} h^{\mu\nu} (\partial_\mu \mathcal{A}_\alpha \partial_\nu \mathcal{A}^\alpha + \partial_\alpha \mathcal{A}_\mu \partial^\alpha \mathcal{A}_\nu \\ &\quad \left. - \partial_\alpha \mathcal{A}_\mu \partial_\nu \mathcal{A}^\alpha - \partial_\alpha \mathcal{A}_\nu \partial_\mu \mathcal{A}^\alpha) \right\} , \end{aligned} \quad (13)$$

and for the matter Lagrangian

$$\begin{aligned} \mathcal{L}_{\text{m}}^{(1)} &= \frac{1}{2} \kappa h (|\partial_\mu \phi|^2 - m^2 |\phi|^2) - \kappa h^{\mu\nu} (\partial_\mu \phi^* \partial_\nu \phi) \\ &\quad + ie_q \kappa (\mathcal{A}_\mu \partial^\mu \phi^* \phi - \mathcal{A}_\mu \phi^* \partial^\mu \phi) + e_q^2 \mathcal{A}_\mu \mathcal{A}^\mu |\phi|^2 \\ &\quad + \frac{1}{2} \kappa h [\partial_\mu \phi^* (ie_q \mathcal{A}^\mu) \phi - (ie_q \mathcal{A}^\mu) \phi^* \partial_\mu \phi] \\ &\quad - \kappa h^{\mu\nu} [\partial_\mu \phi^* (ie_q \mathcal{A}_\nu) \phi + (ie_q \mathcal{A}_\mu) \phi^* \partial_\nu \phi] \end{aligned} \quad (14)$$

The Feynman rules for interaction vertices between photons, complex scalars and gravitons can be derived from (13) and (14), [14], and are presented in Appendix A.

III. GENERAL RELATIVITY AND SCALAR QED AS AN EFFECTIVE FIELD THEORY

The treatment of general relativity as a quantum theory has been discussed in several publications, e.g. [1, 17, 18, 19, 20, 21, 22]. The standard formalism would start from the action (2) and consider the metric as the gravitational field which has to be quantized in the usual way. However, such a theory is not self-contained, since at each order in perturbation theory loop diagrams generate new terms which are not present in the original action [1, 2, 3, 23]. Because of this renormalization problem, a consistent definition of general relativity as a *fundamental* quantum theory is unknown so far [11].

As mentioned above, the problem can be solved by considering general relativity as an effective quantum field theory at low energies [7, 8, 9]. In this case, the effective Lagrangian for pure gravity (without a cosmological constant) has the expansion

$$\mathcal{L}_{\text{grav}} = \sqrt{-g} \left\{ \frac{2\mathcal{R}}{\kappa^2} + a_1 \mathcal{R}^2 + a_2 \mathcal{R}_{\mu\nu} \mathcal{R}^{\mu\nu} + \dots \right\} . \quad (15)$$

Here, the coefficients a_1 and a_2 are energy scale dependent coupling constants in the effective theory which –

in principle – could be determined by an experimental measurement. The ellipses denote higher-derivative couplings which are even more suppressed at low energies. Similarly, as shown in [14] general relativity and scalar QED can be described in the effective-theory framework as well. The 1-loop singularities from graviton and photon exchange can be absorbed into the Lagrangian

$$\Delta\mathcal{L}_{\text{photon}} = \sqrt{-g} \left\{ c_1 T_{\mu\nu}^2 + c_2 \mathcal{R}_{\mu\nu} T^{\mu\nu} + \dots \right\}, \quad (16)$$

where $T_{\mu\nu} = \mathcal{F}_{\mu\alpha}\mathcal{F}_{\nu}^{\alpha} - \frac{1}{4}g_{\mu\nu}\mathcal{F}_{\alpha\beta}\mathcal{F}^{\alpha\beta}$ is the Maxwell stress tensor [4, 5, 14]. For the matter Lagrangian describing the complex scalar fields, one obtains additional terms [14]

$$\mathcal{L}_{\text{scalar}} = \sqrt{-g} \left\{ d_1 \mathcal{R}^{\mu\nu} \partial_{\mu} \phi^* \partial_{\nu} \phi + d_2 \mathcal{R} |\partial_{\mu} \phi|^2 + d_3 \mathcal{R}^2 |\phi|^2 + \dots \right\} \quad (17)$$

An explicit discussion of the coefficients $a_1, a_2, c_1, c_2, d_1, d_2$ and d_3 , can be found in [7, 8, 9, 13] and [14].

IV. SCATTERING MATRIX AND POTENTIAL

According to [7, 8, 9, 13, 14], the scattering amplitude for two massive, charged scalar particles as a function of the momentum transfer $q^2 = (k - k')^2 = (p - p')^2$ can be expanded as

$$\mathcal{M}(q) \sim \left[A + Bq^2 + \dots + (\alpha_1 \kappa^2 + \alpha_2 e^2) \frac{1}{q^2} + \beta_1 e^2 \kappa^2 \log(-q^2) + \beta_2 e^2 \kappa^2 \frac{1}{\sqrt{-q^2}} + \dots \right], \quad (18)$$

where the coefficients $A, B, \dots$ and α_i, β_i depend on the particle masses m_i and charges e_i/e . The terms with $A, B, \dots$ in this equation are analytical in q^2 and correspond to local interactions. They are relevant for large momentum transfer and are also related to the UV renormalization of the theory, but will not be of further interest in this work. Rather, we concentrate on the nonanalytical terms, parametrized by the coefficients α_i and β_i , which will induce the nonlocal, long-ranged interactions described by the nonrelativistic potential. The nonanalytical parts in the scattering-matrix also ensure unitarity [7, 8, 9, 13, 14].

The nonrelativistic potential will be calculated from the scattering amplitude via the spatial Fourier transform [7, 8, 9, 10, 13, 14]

$$V(r) = -\frac{1}{2m_1} \frac{1}{2m_2} \int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{r}} \mathcal{M}(\vec{q}). \quad (19)$$

The non-analytical terms in $\mathcal{M}$ correspond to an expansion in powers of $\frac{1}{r}$ for $V(r)$. For alternative definitions of the potential, see the discussion in, e.g., [24, 25, 26].




FIG. 1: The two tree diagrams.

V. RESULTS FOR THE FEYNMAN DIAGRAMS

In the following section we discuss the calculation of the various Feynman diagrams and their contribution to the scattering amplitude. At the end of this section, we summarize the one-loop result for the nonrelativistic potential which represents the main result of this paper. Part of the calculation for the Feynman diagrams has been done with the help of Mathematica 5.2TM[31] using the Ricci-application©[32] for tensor calculus. The Fourier transformations, necessary to obtain the potential in coordinate space, are listed in Appendix C.

A. Tree Diagrams

The tree diagrams contributing to the scattering amplitude are shown in Fig. 1. The (incoming/outgoing) momenta for the first particle are denoted as (k/k') with the (mass/charge) being (m_1/e_1) , and for the second particle as (p/p') and (m_2/e_2) , respectively. They have the form

$$i\mathcal{M}_{1(a)}(q) = \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{q^2} \right] \tau_1^{\alpha\beta}(p, p', m_2) \quad (20)$$

and

$$i\mathcal{M}_{1(b)}(q) = \tau_2^{\gamma}(k, k', e_1) \left[\frac{-i\eta_{\gamma\delta}}{q^2} \right] \tau_2^{\delta}(p, p', e_2). \quad (21)$$

where the quantities τ_i and $\mathcal{P}_{\mu\nu\alpha\beta}$ are given in Appendix A. After Fourier transformation we recover the Newtonian nonrelativistic potential

$$V_{1(a)}(r) = -G \frac{m_1 m_2}{r} \quad (22)$$

and the classical Coulomb potential

$$V_{1(b)}(r) = \frac{1}{4\pi} \frac{e_1 e_2}{r}. \quad (23)$$

B. Vertex Corrections

There are two classes of vertex corrections, involving gravitational effects only, as shown in Fig. 2. The contri-




FIG. 2: Vertex corrections from purely gravitational effects. (a) and (b) contain the $\phi^2 h$ coupling. (c) and (d) the $\phi^2 h^2$ coupling.

butions from Fig. 2(a) have the form

$$\begin{aligned}
 i\mathcal{M}_{2(a)}(q) = & \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{q^2} \right] \int \frac{d^4\ell}{(2\pi)^4} \\
 & \cdot \tau_7^{\alpha\beta\rho\sigma\gamma\delta}(\ell, -q) \left[\frac{i\mathcal{P}_{\rho\sigma\lambda\epsilon}}{\ell^2} \right] \tau_1^{\lambda\epsilon}(p, p - \ell, m_2) \\
 & \cdot \left[\frac{i}{(\ell - p)^2 - m_2^2} \right] \tau_1^{\tau\phi}(p - \ell, p', m_2) \\
 & \cdot \left[\frac{i\mathcal{P}_{\gamma\delta\tau\phi}}{(\ell + q)^2} \right], \quad (24)
 \end{aligned}$$

and for Fig. 2(b)

$$\begin{aligned}
 i\mathcal{M}_{2(b)}(q) = & \tau_1^{\mu\nu}(p, p', m_2) \left[\frac{i\mathcal{P}_{\alpha\beta\mu\nu}}{q^2} \right] \int \frac{d^4\ell}{(2\pi)^2} \\
 & \cdot \tau_1^{\lambda\epsilon}(k, k + \ell, m_1) \left[\frac{i\mathcal{P}_{\lambda\epsilon\rho\sigma}}{\ell^2} \right] \tau_7^{\alpha\beta\rho\sigma\gamma\delta}(-\ell, q) \\
 & \cdot \left[\frac{i\mathcal{P}_{\gamma\delta\tau\phi}}{(\ell + q)^2} \right] \tau_1^{\tau\phi}(\ell + k, k', m_1) \\
 & \cdot \left[\frac{i}{(\ell + k)^2 - m_1^2} \right]. \quad (25)
 \end{aligned}$$

For the graviton loop diagrams, Fig. 2(c) and (d), we find

$$\begin{aligned}
 i\mathcal{M}_{2(c)}(q) = & \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{q^2} \right] \frac{1}{2!} \int \frac{d^4\ell}{(2\pi)^4} \\
 & \cdot \tau_7^{\alpha\beta\rho\sigma\gamma\delta}(\ell, -q) \left[\frac{i\mathcal{P}_{\rho\sigma\lambda\epsilon}}{\ell^2} \right] \tau_3^{\tau\phi\lambda\epsilon}(p, p', m_2) \\
 & \cdot \left[\frac{i\mathcal{P}_{\tau\phi\gamma\delta}}{(\ell + q)^2} \right], \quad (26)
 \end{aligned}$$

$$\begin{aligned}
 i\mathcal{M}_{2(d)}(q) = & \tau_1^{\mu\nu}(p, p', m_2) \left[\frac{i\mathcal{P}_{\alpha\beta\mu\nu}}{q^2} \right] \frac{1}{2!} \int \frac{d^4\ell}{(2\pi)^2} \\
 & \cdot \tau_3^{\tau\phi\lambda\epsilon}(k, k', m_1) \left[\frac{i\mathcal{P}_{\lambda\epsilon\rho\sigma}}{\ell^2} \right] \tau_7^{\alpha\beta\rho\sigma\gamma\delta}(-\ell, q) \\
 & \cdot \left[\frac{i\mathcal{P}_{\gamma\delta\tau\phi}}{(\ell + q)^2} \right]. \quad (27)
 \end{aligned}$$

Again, the various functions τ_i can be found in the Appendix. Contracting the Lorentz indices and performing the loop integration, we find

$$\begin{aligned}
 i\mathcal{M}_{2(a+b)}(q) = & -8iG^2m_1^2m_2^2 \left[\frac{\pi^2(m_1 + m_2)}{\sqrt{-q^2}} + \frac{5}{3} \log(-q^2) \right], \quad (28)
 \end{aligned}$$

$$i\mathcal{M}_{2(c+d)}(q) = \frac{208i}{3} G^2m_1^2m_2^2 \log(-q^2). \quad (29)$$

The Fourier transform yields a contribution to the non-relativistic potential

$$\begin{aligned}
 V_{2(a+b)}(r) = & \frac{G^2m_1m_2(m_1 + m_2)}{r^2} \\
 & - \frac{5}{3} \frac{G^2m_1m_2}{\pi r^3}, \quad (30)
 \end{aligned}$$

$$V_{2(c+d)}(r) = \frac{26}{3} \frac{G^2m_1m_2}{\pi r^3}. \quad (31)$$

Including electromagnetic interactions in SQED we obtain additional vertex corrections, shown in Figs. 3, 4. The results for the scattering amplitudes from the diagrams in Figs. 3(a) – (d) are given by

$$\begin{aligned}
 i\mathcal{M}_{3(a+b)}(q) = & -\frac{5i}{2} G(e_1^2m_2^2 + e_2^2m_1^2) \log(-q^2) \\
 & - iG(e_2^2m_1^2m_2 + e_1^2m_2^2m_1) \frac{\pi}{\sqrt{-q^2}}, \quad (32)
 \end{aligned}$$

$$i\mathcal{M}_{3(c+d)}(q) = -\frac{i}{6} G(e_1^2m_2^2 + e_2^2m_1^2) \log(-q^2), \quad (33)$$




FIG. 3: Vertex corrections involving gravitational and electromagnetic interactions (graviton 1-particle reducible).




FIG. 4: Vertex corrections involving gravitational and electromagnetic interactions (photon 1-particle reducible).

which contributes to the nonrelativistic potential as

$$V_{3(a-d)}(r) = \frac{G(m_1 e_2^2 + m_2 e_1^2)}{8\pi^2 r^2} - \frac{G(\frac{m_2}{m_1} e_1^2 + \frac{m_1}{m_2} e_2^2)}{3\pi r^3}. \quad (34)$$

Note, that the diagrams Figs. 3(c) and (d) are associated with a symmetry factor 1/2. Further details of the computation can be found in Appendix D. Similarly, the diagrams in Fig. 4 yield

$$i\mathcal{M}_{4(a-f)}(q) = 2iGm_1m_2e_1e_2 \frac{\pi(m_1 + m_2)}{\sqrt{-q^2}}, \quad (35)$$

and

$$V_{4(a-f)}(r) = -\frac{Ge_1e_2(m_1 + m_2)}{4\pi r^2}. \quad (36)$$

C. Triangle Diagrams

The triangle diagrams in Fig. 5 arise from purely gravitational interactions. From the expressions in Appendix D we compute

$$i\mathcal{M}_{5(a+b)}(q) = 8iG^2m_1^2m_2^2 \left[28 \log(-q^2) + \frac{4\pi^2(m_1 + m_2)}{\sqrt{-q^2}} \right] \quad (37)$$




FIG. 5: Triangle diagrams with only gravitational interactions.




FIG. 6: Triangle diagrams with gravitational and electromagnetic interactions.

which contributes to the potential as

$$V_{5(a+b)}(r) = -\frac{G^2m_1m_2}{r} \left[\frac{4(m_1 + m_2)}{r} - \frac{28}{\pi r^2} \right]. \quad (38)$$

The additional triangle diagrams in SQED are shown in Fig. 6. Their contribution to the scattering amplitude reads

$$i\mathcal{M}_{6(a-d)}(q) = 4im_1m_2 \frac{Ge_1e_2}{\pi} \left[-8 \log(-q^2) - \frac{2\pi^2(m_1 + m_2)}{\sqrt{-q^2}} \right], \quad (39)$$

resulting in the potential

$$V_{6(a-d)}(r) = \frac{Ge_1e_2(m_1 + m_2)}{\pi r^2} - \frac{4Ge_1e_2}{\pi^2 r^3}. \quad (40)$$




FIG. 7: Box and crossed box diagrams from purely gravitational interactions.

D. Box and Crossed Box Diagrams

The contributions of the box and crossed box diagrams in Fig. 7 read

$$i\mathcal{M}_{7(a)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_1^{\mu\nu}(k, k+\ell, m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{\ell^2} \right] \cdot \tau_1^{\alpha\beta}(p, p-\ell, m_2) \left[\frac{i}{(\ell-p)^2 - m_2^2} \right] \cdot \tau_1^{\gamma\delta}(p-\ell, p', m_2) \left[\frac{i\mathcal{P}_{\gamma\delta\rho\sigma}}{(\ell+q)^2} \right] \cdot \tau_1^{\rho\sigma}(k+\ell, k', m_1) \left[\frac{i}{(\ell+k)^2 - m_1^2} \right], \quad (41)$$

and

$$i\mathcal{M}_{7(b)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_1^{\mu\nu}(k, \ell+k, m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{\ell^2} \right] \cdot \tau_1^{\alpha\beta}(\ell+p', p', m_2) \left[\frac{i}{(\ell+p')^2 - m_2^2} \right] \cdot \tau_1^{\gamma\delta}(p, \ell+p', m_2) \left[\frac{i\mathcal{P}_{\gamma\delta\rho\sigma}}{(\ell+q)^2} \right] \cdot \tau_1^{\rho\sigma}(k+\ell, k', m_1) \left[\frac{i}{(\ell+k)^2 - m_1^2} \right]. \quad (42)$$

Again, we are only interested in the nonanalytic terms, and one may exploit some simplifications which reduce parts of the amplitude to three- and two-point functions, see Appendix B,

$$i\mathcal{M}_{7(a+b)}^{\text{red}}(q) = -64 i G^2 m_1^2 m_2^2 \log(-q^2). \quad (43)$$

For the remaining irreducible parts we find

$$i\mathcal{M}_{7(a+b)}^{\text{irred}}(q) = -\frac{184}{3} i G^2 m_1^2 m_2^2 \log(-q^2). \quad (44)$$

In the nonrelativistic limit this combines to the potential

$$V_{7(a+b)} = -\frac{47}{3} \frac{G^2 m_1 m_2}{\pi r^3}. \quad (45)$$

Similarly, including electromagnetic interactions, the box and crossed box diagrams in Fig. 8 give a reducible part,

$$i\mathcal{M}_{8(a-d)}^{\text{red}}(q) = 32 i G m_1 m_2 \frac{e_1 e_2}{4\pi} \log(-q^2), \quad (46)$$




FIG. 8: Box and crossed box diagrams with gravitational and electromagnetic interactions.

and an irreducible part,

$$i\mathcal{M}_{8(a-d)}^{\text{irred}}(q) = \frac{224}{3} i G m_1 m_2 \frac{e_1 e_2}{4\pi} \log(-q^2) \quad (47)$$

The contribution to the potential is

$$V_{8(a-d)}(r) = \frac{10}{3} \frac{G e_1 e_2}{\pi^2 r^3}. \quad (48)$$

E. Circular Diagram

The calculation of the circular diagrams in Fig. 9 is straightforward. We find the nonanalytic contributions to the scattering amplitude

$$i\mathcal{M}_{9(a)}(q) = -176 i G^2 m_1^2 m_2^2 \log(-q^2) \quad (49)$$

and

$$i\mathcal{M}_{9(b)}(q) = \frac{16}{\pi} i G m_1 m_2 e_1 e_2 \log(-q^2), \quad (50)$$




FIG. 9: Circular diagrams: (a) purely gravitational; (b) gravitational and electromagnetic interactions.




FIG. 10: The set of vacuum polarization diagrams which contribute to the potential.

and the resulting contribution to the potential

$$V_{9(a)}(r) = -22 \frac{G^2 m_1 m_2}{\pi r^3}, \quad (51)$$

and

$$V_{9(b)}(r) = 2 \frac{G e_1 e_2}{\pi^2 r^3}. \quad (52)$$

F. Vacuum Polarization Diagram

Treating general relativity as an effective field theory, we have to deal with the vacuum polarization diagrams Fig. 10(a)–(d) as well. The graviton self-energy is obtained from the subdiagrams in Fig. 10(a-b). Its contribution to the effective Lagrangian has been worked out by 't Hooft and Veltman [1, 2, 7, 8],

$$\mathcal{L} = \frac{-1}{16\pi^2} \left(\frac{1}{120} \bar{\mathcal{R}}^2 + \frac{7}{20} \mathcal{R}_{\mu\nu} \mathcal{R}^{\mu\nu} \right) \log(-q^2). \quad (53)$$

From this Lagrangian, one obtains the vacuum polarization tensor

$$\begin{aligned} \tilde{\Pi}_{\alpha\beta,\gamma\delta} = \frac{-2G}{\pi} & \left[\frac{21}{120} q^4 \mathbb{1}_{\alpha\beta\gamma\delta} + \frac{23}{120} q^4 \eta_{\alpha\beta} \eta_{\gamma\delta} \right. \\ & - \frac{23}{120} (\eta_{\alpha\beta} q_\gamma q_\delta + \eta_{\gamma\delta} q_\alpha q_\beta) q^2 \\ & - \frac{21}{240} (q_\alpha q_\delta \eta_{\beta\gamma} + q_\beta q_\delta \eta_{\alpha\gamma} \\ & + q_\alpha q_\gamma \eta_{\beta\delta} + q_\beta q_\gamma \eta_{\alpha\delta}) q^2 \\ & \left. + \frac{11}{30} q_\alpha q_\beta q_\gamma q_\delta \right] \log(-q^2). \quad (54) \end{aligned}$$

Inserting this into the scattering amplitudes in Fig. 10(a+b), one obtains

$$\begin{aligned} i\mathcal{M}_{10(a+b)}(q) &= \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{q^2} \right] \\ &\quad \cdot \tilde{\Pi}^{\alpha\beta,\gamma\delta}(q) \left[\frac{i\mathcal{P}_{\gamma\delta\rho\sigma}}{q^2} \right] \tau_1^{\rho\sigma}(p, p', m_2), \\ &= -\frac{172i}{15} G^2 m_1^2 m_2^2 \log(-q^2), \quad (55) \end{aligned}$$

which yields the potential

$$V_{10(a+b)}(r) = -\frac{43}{30} \frac{G^2 m_1 m_2}{\pi r^3}. \quad (56)$$

Furthermore we have to treat the scalar and photon contribution to the graviton self-energy in Figs. 10(c) and 10(d), respectively. The contribution of massive scalar particles only contains analytical terms and is therefore irrelevant for the calculation of the nonrelativistic potential. If we have N additional massless (uncharged) scalars in the theory, they would contribute to the graviton polarization tensor as

$$\begin{aligned} \bar{\Pi}_{\alpha\beta,\gamma\delta} = N \frac{-G}{\pi} & \left[\frac{1}{20} (q_\alpha q_\beta - q^2 \eta_{\alpha\beta}) (q_\gamma q_\delta - q^2 \eta_{\gamma\delta}) \right. \\ & + \frac{1}{30} (q_\alpha q_\gamma - q^2 \eta_{\alpha\gamma}) (q_\beta q_\delta - q^2 \eta_{\beta\delta}) \\ & \left. + \frac{1}{30} (q_\alpha q_\delta - q^2 \eta_{\alpha\delta}) (q_\beta q_\gamma - q^2 \eta_{\beta\gamma}) \right] \log(-q^2). \quad (57) \end{aligned}$$

Inserting this into the scattering amplitude yields

$$i\mathcal{M}_{10(c)}(q) = -N \frac{8i}{5} G^2 m_1^2 m_2^2 \log(-q^2), \quad (58)$$

which corresponds to a contribution to the nonrelativistic potential

$$V_{10(c)}(r) = \frac{-N}{5} \frac{G^2 m_1 m_2}{\pi r^3}. \quad (59)$$

Finally, one calculates the photon-loop contribution to the polarization tensor

$$\begin{aligned} \hat{\Pi}_{\alpha\beta,\gamma\delta} = \frac{G}{\pi} & \left[\frac{1}{15} (q_\alpha q_\beta - q^2 \eta_{\alpha\beta}) (q_\gamma q_\delta - q^2 \eta_{\gamma\delta}) \right. \\ & - \frac{1}{10} (q_\alpha q_\gamma - q^2 \eta_{\alpha\gamma}) (q_\beta q_\delta - q^2 \eta_{\beta\delta}) \\ & \left. - \frac{1}{10} (q_\alpha q_\delta - q^2 \eta_{\alpha\delta}) (q_\beta q_\gamma - q^2 \eta_{\beta\gamma}) \right] \log(-q^2), \quad (60) \end{aligned}$$

which yields

$$i\mathcal{M}_{10(d)}(q) = -\frac{64i}{15} G^2 m_1^2 m_2^2 \log(-q^2), \quad (61)$$




FIG. 11: The one-loop contribution to the photon self-energy in SQED.

and

$$V_{10(d)} = -\frac{4}{15} \frac{G^2 m_1 m_2}{\pi r^2}. \quad (62)$$

The photon self-energy in SQED only receives a one-loop contribution from the mixed vacuum polarization diagram in Fig. 11,

$$\Pi^{\gamma\tau}(q) = \frac{iG}{\pi} \left(\frac{q^2}{3} \right) (q^\gamma q^\tau - q^2 \eta^{\gamma\tau}) \log(-q^2), \quad (63)$$

which yields

$$i\mathcal{M}_{11}(q) = \frac{4i}{3\pi} G m_1 m_2 e_1 e_2 \log(-q^2), \quad (64)$$

and

$$V_{11}(r) = \frac{1}{6} \frac{G e_1 e_2}{\pi^2 r^3}. \quad (65)$$

We note that the one-loop diagrams in Fig. 12 only contribute analytical terms to the scattering amplitude and thus do not contribute to the nonrelativistic potential, see also [7, 14].

G. Final result for the nonrelativistic potential

In this paragraph we are going to summarize the main result of our paper: the nonrelativistic potential which is obtained from combining the contributions of the various diagrams discussed above. Let us first concentrate on the purely gravitational effects, for which we obtain

$$V_{\text{GR}}(r) = -\frac{G m_1 m_2}{r} \left[1 + 3 \frac{G(m_1 + m_2)}{c^2 r} + \frac{131 + 6N}{30\pi} \frac{G\hbar}{c^3 r^2} \right]. \quad (66)$$

Here, we have restored the dependence on c and $\hbar$ to distinguish the effects from general relativity and quantum corrections. We recall that N denotes the number of massless (uncharged) scalars, e.g., “scalar neutrinos.” When comparing our result (66) with [13], we have to take into account that in [13] the quantum corrections to the graviton propagator from massless scalar and photon




FIG. 12: The set of diagrams which only give analytical contributions to the scattering amplitude.

loops have not been considered. Apart from this, we fully agree with [13]. The discrepancies between [13, 27] and the original calculations in [7, 8, 9] and the results of [12] have already been discussed, and we thus confirm the conclusions of [13]. Concerning the contributions from massless scalars to the polarization tensor in (57), we disagree with a previous calculation in [10]. The photon contribution in (60) coincides with the result in [28]. Including electromagnetic effects in scalar QED, we find

$$V_{\text{SQED}}(r) = V_{\text{GR}}(r) + \frac{\tilde{\alpha} \mathbf{e}_1 \mathbf{e}_2}{r} \left(1 + 3 \frac{G(m_1 + m_2)}{c^2 r} + \frac{6}{\pi} \frac{G\hbar}{c^3 r^2} \right) + \frac{1}{2} \frac{(m_1 \mathbf{e}_2^2 + m_2 \mathbf{e}_1^2) G \tilde{\alpha}}{c^2 r^2} - \frac{4}{3\pi} \left(\frac{m_2^2 \mathbf{e}_1^2 + m_1^2 \mathbf{e}_2^2}{m_1 m_2} \right) \frac{G \tilde{\alpha} \hbar}{c^3 r^3}. \quad (67)$$

Here we have introduced $\tilde{\alpha} = \hbar c/137$ and the charges $\mathbf{e}_1$ and $\mathbf{e}_2$ are normalized in units of the elementary charge. Again, our result for the potential (67) is in agreement with [14, 29]. [A calculational error in the expression for the box diagrams (47) quoted in a previous preprint version of this paper [30] has been corrected.]

VI. SUMMARY

We have calculated the one-loop corrections to the Newtonian and Coulomb potential for massive and electrically charged scalar particles, treating general relativity as an effective quantum field theory. The nonrelativistic potential is obtained from the scattering amplitude by considering the nonanalytic terms in the momentum transfer q^2 . After Fourier transformation, we identify relativistic corrections of order $Gm/c^2 r$ and genuine quantum corrections of order $G\hbar/c^3 r^2$. Our results for the leading quantum corrections to the Newtonian and Coulomb laws are in agreement with [13] and [14, 29]. Concerning the quantum corrections to the graviton propagator, we reproduced the contribution from the photon loop in [28], but differ for the case of massless scalar loops in [10]. It is straight forward to generalize the effective-theory formalism to, for instance, the case of massive charged spin- $\frac{1}{2}$ fermions which has been discussed in [15, 16]. In this way one could unambiguously predict the

quantum-gravitational corrections to low-energy observables within the Standard Model. However, one has to be aware that the size of the calculated corrections is – of course – extremely small, $G\hbar/c^3 r^2 = 2.6 \cdot 10^{-40} \text{ fm}^2/r^2$, and it will thus be practically impossible to verify the theoretical results by experimental measurements in the foreseeable future [9].

Acknowledgments

I would like to thank my supervisors Th. Mannel and A. Khodjamirian for the encouragement and many helpful comments. Special thanks go to N. E. J. Bjerrum-Bohr, University of Wales, Swansea, and Institute for Advanced Study, Princeton, 2006-2007, for discussions on this topic, and to Th. Feldmann for critical discussions and comments on this manuscript.

APPENDIX A: FEYNMAN RULES

1. Propagators

The propagator for a massive scalar field is given by

$$\text{---}\overset{q}{\longrightarrow}\text{---} = \frac{i}{q^2 - m^2 + i\epsilon}.$$

The photon propagator in harmonic (Feynman) gauge is given by

$$\mu \text{ ~~~~~ } \nu = \frac{-i\eta^{\mu\nu}}{q^2 + i\epsilon}.$$

In harmonic gauge the graviton propagator has been discussed in [1, 7]. An explicit derivation can be found in [27] with the result

$$\mu\nu \text{ ~~~~~ } \alpha\beta = \frac{i\mathcal{P}^{\mu\nu\alpha\beta}}{q^2 + i\epsilon}$$

where the polarization sum is given by

$$\mathcal{P}^{\mu\nu\alpha\beta} = \frac{1}{2}(\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\mu\beta}\eta^{\nu\alpha} - \eta^{\mu\nu}\eta^{\alpha\beta}).$$

2. Vertices

The 1-graviton-2-scalar vertex is given by

$$\mu\nu \text{ ~~~~~ } \begin{array}{c} p' \\ \nearrow \\ p \end{array} = \tau_1^{\mu\nu}(p, p', m),$$

with

$$\tau_1^{\mu\nu}(p, p', m) = \frac{-i\kappa}{2}\{p^\mu p'^\nu + p^\nu p'^\mu - \eta^{\mu\nu}[(p \cdot p') - m^2]\}$$

The 1-photon-2-scalar vertex is found to be

$$\begin{array}{c} \gamma \text{ ~~~~~ } \begin{array}{c} p' \\ \nearrow \\ p \end{array} \\ \overrightarrow{q} \end{array} = \tau_2^\gamma(p, p', e) = -ie(p^\gamma + p'^\gamma).$$

The 2-graviton-2-scalar-vertex reads

$$\begin{array}{c} \eta\lambda \text{ ~~~~~ } \begin{array}{c} p' \\ \nearrow \\ p \end{array} \\ \rho\sigma \end{array} = \tau_3^{\eta\lambda\rho\sigma}(p, p', m)$$

with

$$\begin{aligned} \tau_3^{\eta\lambda\rho\sigma}(p, p', m) = i\kappa^2 \Big\{ & \left[\mathbb{1}^{\eta\lambda\alpha\delta} \mathbb{1}^{\rho\sigma\beta}{}_\delta - \frac{1}{4} \left(\eta^{\eta\lambda} \mathbb{1}^{\rho\sigma\alpha\beta} \right. \right. \\ & \left. \left. + \eta^{\rho\sigma} \mathbb{1}^{\eta\lambda\alpha\beta} \right) \right] (p_\alpha p'_\beta + p_\beta p'_\alpha) - \frac{1}{2} \left[\mathbb{1}^{\eta\lambda\rho\sigma} \right. \\ & \left. - \frac{1}{2} \eta^{\eta\lambda} \eta^{\rho\sigma} \right] \left((p \cdot p') - m^2 \right) \Big\} \end{aligned}$$

and

$$\mathbb{1}_{\alpha\beta\gamma\delta} = \frac{1}{2}(\eta_{\alpha\gamma}\eta_{\beta\delta} + \eta_{\alpha\delta}\eta_{\beta\gamma}).$$

The 2-photon-1-graviton vertex is given by

$$\begin{array}{c} \delta \\ \nearrow \\ \mu\nu \text{ ~~~~~ } \begin{array}{c} p' \\ \nearrow \\ p \end{array} \\ \overrightarrow{q} \end{array} = \tau_4^{\mu\nu(\gamma\delta)}(p, p'),$$

with


$$\begin{aligned} \tau_4^{\mu\nu(\gamma\delta)}(p, p') = i\kappa \Big\{ & \mathcal{P}^{\mu\nu(\gamma\delta)}(p \cdot p') + \frac{1}{2} \left[\eta^{\mu\nu} p^\delta p'^\gamma \right. \\ & + \eta^{\gamma\delta} (p^\mu p'^\nu + p^\nu p'^\mu) - (\eta^{\mu\delta} p'^\gamma p^\nu + \eta^{\nu\delta} p'^\gamma p^\mu \\ & \left. + \eta^{\nu\gamma} p'^\mu p^\delta + \eta^{\mu\gamma} p'^\nu p^\delta) \right] \Big\}. \end{aligned}$$

The 2-scalar-2-photon vertex in scalar QED reads


$$\begin{array}{c} \delta \\ \nearrow \\ \gamma \text{ ~~~~~ } \begin{array}{c} p' \\ \nearrow \\ p \end{array} \\ \gamma \end{array} = \tau_5^{\gamma\delta}(p, p', e) = 2 i e^2 \eta^{\gamma\delta}.$$

The 2-scalar-1-photon-1-graviton is found to be

is found as [7, 8, 14]



$$= \tau_6^{\mu\nu(\gamma)}(p, p', e)$$



$$= \tau_7^{\mu\nu}{}_{\alpha\beta\gamma\delta}(k, q),$$

with

$$\tau_6^{\mu\nu(\gamma)}(p, p', e) = i e \kappa [\mathcal{P}^{\mu\nu\alpha\gamma}(p + p')_\alpha].$$

Using the background field method the 3-graviton vertex with

$$\begin{aligned} \tau_7^{\mu\nu}{}_{\alpha\beta\gamma\delta} = & -\frac{i\kappa}{2} \cdot \left\{ \mathcal{P}_{\alpha\beta\gamma\delta} \left[k^\mu k^\nu + (k - q)^\mu (k - q)^\nu + q^\mu q^\nu - \frac{3}{2} \eta^{\mu\nu} q^2 \right] + 2q_\lambda q_\sigma [\mathbb{1}_{\alpha\beta}{}^{\sigma\lambda} \mathbb{1}_{\gamma\delta}{}^{\mu\nu} + \mathbb{1}_{\gamma\delta}{}^{\sigma\lambda} \mathbb{1}_{\alpha\beta}{}^{\mu\nu} - \mathbb{1}_{\alpha\beta}{}^{\mu\sigma} \mathbb{1}_{\gamma\delta}{}^{\nu\lambda} \right. \\ & - \mathbb{1}_{\gamma\delta}{}^{\mu\sigma} \mathbb{1}_{\alpha\beta}{}^{\nu\lambda}] + [q_\lambda q^\mu (\eta_{\alpha\beta} \mathbb{1}_{\gamma\delta}{}^{\nu\lambda} + \eta_{\gamma\delta} \mathbb{1}_{\alpha\beta}{}^{\nu\lambda}) + q_\lambda q^\nu (\eta_{\alpha\beta} \mathbb{1}_{\gamma\delta}{}^{\mu\lambda} + \eta_{\gamma\delta} \mathbb{1}_{\alpha\beta}{}^{\mu\lambda}) \\ & - q^2 (\eta_{\alpha\beta} \mathbb{1}_{\gamma\delta}{}^{\mu\nu} + \eta_{\gamma\delta} \mathbb{1}_{\alpha\beta}{}^{\mu\nu}) - \eta^{\mu\nu} q_\lambda q_\sigma (\eta_{\alpha\beta} \mathbb{1}_{\gamma\delta}{}^{\sigma\lambda} + \eta_{\gamma\delta} \mathbb{1}_{\alpha\beta}{}^{\sigma\lambda})] \\ & + [2q_\lambda \{ \mathbb{1}_{\alpha\beta}{}^{\lambda\sigma} \mathbb{1}_{\gamma\delta\sigma}{}^\nu (k - q)^\mu + \mathbb{1}_{\alpha\beta}{}^{\lambda\sigma} \mathbb{1}_{\gamma\delta\sigma}{}^\mu (k - q)^\nu - \mathbb{1}_{\gamma\delta}{}^{\lambda\sigma} \mathbb{1}_{\alpha\beta\sigma}{}^\nu k^\mu - \mathbb{1}_{\gamma\delta}{}^{\lambda\sigma} \mathbb{1}_{\alpha\beta\sigma}{}^\mu k^\nu \} \\ & + q^2 (\mathbb{1}_{\alpha\beta\sigma}{}^\mu \mathbb{1}_{\gamma\delta}{}^{\nu\sigma} + \mathbb{1}_{\alpha\beta}{}^{\nu\sigma} \mathbb{1}_{\gamma\delta\sigma}{}^\mu) + \eta^{\mu\nu} q_\sigma q_\lambda (\mathbb{1}_{\alpha\beta}{}^{\lambda\rho} \mathbb{1}_{\gamma\delta\rho}{}^\sigma + \mathbb{1}_{\gamma\delta}{}^{\lambda\rho} \mathbb{1}_{\alpha\beta\rho}{}^\sigma)] + \left\{ (k^2 + (k - q)^2) \right. \\ & \cdot \left[\mathbb{1}_{\alpha\beta}{}^{\mu\sigma} \mathbb{1}_{\gamma\delta\sigma}{}^\nu + \mathbb{1}_{\gamma\delta}{}^{\mu\sigma} \mathbb{1}_{\alpha\beta\sigma}{}^\nu - \frac{1}{2} \eta^{\mu\nu} \mathcal{P}_{\alpha\beta\gamma\delta} \right] - (\mathbb{1}_{\gamma\delta}{}^{\mu\nu} \eta_{\alpha\beta} k^2 - \mathbb{1}_{\alpha\beta}{}^{\mu\nu} \eta_{\gamma\delta} (k - q)^2) \left. \right\} \Big\}. \end{aligned}$$

APPENDIX B: BASIC INTEGRALS

For the calculation of the Feynman diagrams we need the following two-point integrals:

$$J = \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell + q)^2} = \frac{i}{32\pi^2} [-2L] + \dots \quad (\text{B1})$$

$$J_\mu = \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu}{\ell^2(\ell + q)^2} = \frac{i}{32\pi^2} [q_\mu L] + \dots \quad (\text{B2})$$

$$J_{\mu\nu} = \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu \ell_\nu}{\ell^2(\ell + q)^2} = \frac{i}{32\pi^2} \left[q_\mu q_\nu \left(-\frac{2}{3} L \right) - q^2 \eta_{\mu\nu} \left(-\frac{1}{6} L \right) \right] + \dots \quad (\text{B3})$$

$$J_{\mu\nu\alpha} = \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu \ell_\nu \ell_\alpha}{\ell^2(\ell + q)^2} = \frac{i}{32\pi^2} \left[q_\mu q_\nu q_\alpha \left(\frac{1}{2} L \right) - q^2 (q_\alpha \eta_{\mu\nu} + q_\mu \eta_{\nu\alpha} + q_\nu \eta_{\mu\alpha}) \left(\frac{1}{12} L \right) \right] + \dots \quad (\text{B4})$$

$$\begin{aligned} J_{\mu\nu\alpha\beta} = & \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu \ell_\nu \ell_\alpha \ell_\beta}{\ell^2(\ell + q)^2} \\ = & \frac{i}{32\pi^2} \left[q_\mu q_\nu q_\alpha q_\beta \left(-\frac{2}{5} L \right) - q^2 (q_\alpha q_\beta \eta_{\mu\nu} + q_\mu q_\beta \eta_{\nu\alpha} + q_\nu q_\alpha \eta_{\mu\beta} + q_\mu q_\beta \eta_{\nu\alpha} + q_\mu q_\alpha \eta_{\nu\beta} \right. \right. \\ & \left. \left. + q_\mu q_\nu \eta_{\alpha\beta}) \left(-\frac{1}{20} L \right) + q^4 (\eta_{\mu\nu} \eta_{\alpha\beta} + \eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha}) \left(-\frac{1}{120} L \right) \right] + \dots \end{aligned} \quad (\text{B5})$$

together with the three-point integrals

$$I = \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell+q)^2[(\ell+k)^2-m^2]} = \frac{i}{32\pi^2 m^2} [-L-S] + \dots \quad (\text{B6})$$

$$I_\mu = \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu}{\ell^2(\ell+q)^2[(\ell+k)^2-m^2]} = \frac{i}{32\pi^2 m^2} \left\{ k_\mu \left[\left(-1 - \frac{1}{2} \frac{q^2}{m^2} \right) L - \frac{1}{4} \frac{q^2}{m^2} S \right] + q_\mu \left(L + \frac{1}{2} S \right) \right\} + \dots \quad (\text{B7})$$

$$\begin{aligned} I_{\mu\nu} &= \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu \ell_\nu}{\ell^2(\ell+q)^2[(\ell+k)^2-m^2]} \\ &= \frac{i}{32\pi^2 m^2} \left\{ q_\mu q_\nu \left(-L + \frac{3}{8} S \right) + k_\mu k_\nu \left(-\frac{1}{2} \frac{q^2}{m^2} L - \frac{1}{8} \frac{q^2}{m^2} S \right) + (q_\mu k_\nu + q_\nu k_\mu) \left[\left(\frac{1}{2} + \frac{1}{2} \frac{q^2}{m^2} \right) L + \frac{3}{16} \frac{q^2}{m^2} S \right] \right. \\ &\quad \left. + q^2 \eta_{\mu\nu} \left(\frac{1}{4} L + \frac{1}{8} S \right) \right\} + \dots \end{aligned} \quad (\text{B8})$$

$$\begin{aligned} I_{\mu\nu\alpha} &= \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell_\mu \ell_\nu \ell_\alpha}{\ell^2(\ell+q)^2[(\ell+k)^2-m^2]} \\ &= \frac{i}{32\pi^2 m^2} \left\{ q_\mu q_\nu q_\alpha \left(L + \frac{5}{16} S \right) + k_\mu k_\nu k_\alpha \left(-\frac{1}{6} \frac{q^2}{m^2} L \right) + (q_\mu k_\nu k_\alpha + q_\nu k_\mu k_\alpha + q_\alpha k_\mu k_\nu) \left(\frac{1}{3} \frac{q^2}{m^2} L + \frac{1}{16} \frac{q^2}{m^2} S \right) \right. \\ &\quad + (q_\mu q_\nu k_\alpha + q_\mu q_\alpha k_\nu + q_\nu q_\alpha k_\mu) \left[\left(-\frac{1}{3} - \frac{1}{2} \frac{q^2}{m^2} \right) L - \frac{5}{32} \frac{q^2}{m^2} S \right] \\ &\quad \left. + (\eta_{\mu\nu} k_\alpha + \eta_{\mu\alpha} k_\nu + \eta_{\nu\alpha} k_\mu) \left(\frac{1}{12} q^2 L \right) + (\eta_{\mu\nu} q_\alpha + \eta_{\mu\alpha} q_\nu + \eta_{\nu\alpha} q_\mu) \left(-\frac{1}{6} q^2 L - \frac{1}{16} q^2 S \right) \right\} + \dots \end{aligned} \quad (\text{B9})$$

where $L = \log(-q^2)$ and $S = \frac{\pi^2 m}{\sqrt{-q^2}}$, and the external momenta satisfy $k^2 = m^2$ and $k \cdot q = q^2/2$. Here, we have only quoted the leading nonanalytical terms. The ellipses denote higher order nonanalytical contributions and analytical terms. We have checked these integrals independently by explicit (manual) calculation and using MATHEMATICA.

1. Reductions of the Four-Point Integrals

For the box diagrams the following four-point integrals are needed:

$$K = \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} = \frac{i}{16\pi^2 m_1 m_2 q^2} \left[\left(1 - \frac{w}{3m_1 m_2} \right) L \right] + \dots \quad (\text{B10})$$

$$K' = \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell+p')^2-m_2^2]} = \frac{i}{16\pi^2 m_1 m_2 q^2} \left[\left(-1 + \frac{W}{3m_1 m_2} \right) L \right] + \dots \quad (\text{B11})$$

In these equations $w = (k \cdot p) - m_1 m_2$ and $W = (k \cdot p') - m_1 m_2$, which satisfy $W - w = k \cdot (p' - p) = (k \cdot q) = q^2/2$, [14, 16].

For on-shell scalar particles one has the following identities:

$$\ell \cdot q = \frac{1}{2} [(\ell+q)^2 - q^2 - \ell^2], \quad \ell \cdot k = \frac{1}{2} \{ [(\ell+k)^2 - m_1^2] - \ell^2 \} \text{ and } \ell \cdot p = -\frac{1}{2} \{ [(\ell-p)^2 - m_2^2] - \ell^2 \} \quad (\text{B12})$$

Since the terms with $(\ell+q)^2$ and ℓ^2 do not contribute to nonanalytical terms [7, 8, 11, 12, 13, 14, 16], we obtain for

the four-point functions the simplifications/reductions

$$\begin{aligned} \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell \cdot q}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} &= \frac{1}{2} \int \frac{d^4\ell}{(2\pi)^4} \frac{(\ell+q)^2-q^2-\ell^2}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} \\ &\longrightarrow -\frac{q^2}{2} \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} = -\frac{q^2}{2} K, \end{aligned} \quad (\text{B13})$$

$$\begin{aligned} \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell \cdot k}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} &= \frac{1}{2} \int \frac{d^4\ell}{(2\pi)^4} \frac{[(\ell+k)^2-m_1^2]-\ell^2}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} \\ &\longrightarrow \frac{1}{2} \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell+q)^2[(\ell-p)^2-m_2^2]} = \frac{1}{2} I(m_2, p) \end{aligned} \quad (\text{B14})$$

$$\begin{aligned} \int \frac{d^4\ell}{(2\pi)^4} \frac{\ell \cdot p}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} &= -\frac{1}{2} \int \frac{d^4\ell}{(2\pi)^4} \frac{[(\ell-p)^2-m_2^2]-\ell^2}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2][(\ell-p)^2-m_2^2]} \\ &\longrightarrow -\frac{1}{2} \int \frac{d^4\ell}{(2\pi)^4} \frac{1}{\ell^2(\ell+q)^2[(\ell+k)^2-m_1^2]} = -\frac{1}{2} I(m_1, k) \end{aligned} \quad (\text{B15})$$

The four-point functions can thus be reduced to three- or two-point functions, which simplifies the calculation significantly [13, 14, 16].

2. Constraints for the Nonanalytical Terms

The following constraints for the nonanalytical terms are useful

$$J_{\mu\nu}\eta^{\mu\nu} = I_{\mu\nu}\eta^{\mu\nu} = I_{\mu\nu\alpha}\eta^{\mu\nu} = 0, \quad (\text{B16})$$

$$\begin{aligned} J_\mu q^\mu &= -\frac{q^2}{2} J, \quad J_{\mu\nu} q^\nu = -\frac{q^2}{2} J, \\ I_\mu q^\mu &= -\frac{q^2}{2} I, \quad I_{\mu\nu} q^\nu = -\frac{q^2}{2} I_\mu, \end{aligned} \quad (\text{B17})$$

$$I_{\mu\nu\alpha} q^\alpha = -\frac{q^2}{2} I_{\mu\nu},$$

$$I_\mu k^\mu = \frac{1}{2} J, \quad I_{\mu\nu} k^\nu = \frac{1}{2} k_\mu, \quad I_{\mu\nu\alpha} k^\alpha = \frac{1}{2} k_{\mu\nu}. \quad (\text{B18})$$

APPENDIX C: FOURIER TRANSFORMATIONS

The basic Fourier integrals read

$$\int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{r}} \frac{1}{|\vec{q}|^2} = \frac{1}{4\pi r}, \quad (\text{C1})$$

$$\int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{r}} \frac{1}{|\vec{q}|} = \frac{1}{2\pi^2 r^2} \quad (\text{C2})$$

and

$$\int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{r}} \log(\vec{q}^2) = -\frac{1}{2\pi r^3}. \quad (\text{C3})$$

APPENDIX D: FORMAL EXPRESSIONS

Here we collect the formal expressions for the scattering amplitudes, following from applying the Feynman rules given above to the various diagrams. We only quote those results which have not already been given in the body of the text:

- Figs 3(a) and (b):

$$\begin{aligned} i\mathcal{M}_{3(a)}(q) &= \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\rho\sigma}}{q^2} \right] \int \frac{d^4\ell}{(2\pi)^4} \left[\frac{-i\eta_{\alpha\gamma}}{\ell^2} \right] \tau_4^{\rho\sigma(\alpha\beta)}(\ell, \ell+q) \tau_2^\gamma(p, p-\ell) \left[\frac{i}{(\ell-p)^2-m_2^2} \right] \\ &\quad \cdot \tau_2^\delta(p-\ell, p', m_2) \left[\frac{-i\eta_{\delta\beta}}{(\ell+q)^2} \right], \end{aligned} \quad (\text{D1})$$

$$i\mathcal{M}_{3(c)}(q) = \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\rho\sigma}}{q^2} \right] \int \frac{d^4\ell}{(2\pi)^4} \tau_4^{\rho\sigma(\alpha\beta)}(\ell, \ell+q) \left[\frac{-i\eta_{\beta\delta}}{(\ell+q)^2} \right] \tau_5^{\delta\gamma}(p, p', m_2) \left[\frac{-i\eta_{\gamma\alpha}}{\ell^2} \right], \quad (\text{D2})$$

- Figs 4(a) and (e):

$$i\mathcal{M}_{4(a)}(q) = \tau_2^\alpha(k, k', m_1) \left[\frac{-i \eta_{\alpha\beta}}{q^2} \right] \int \frac{d^4\ell}{(2\pi)^4} \tau_4^{\rho\sigma(\beta\gamma)}(q, \ell + q) \cdot \left[\frac{i\mathcal{P}_{\rho\sigma\mu\nu}}{\ell^2} \right] \tau_1^{\mu\nu}(p, p - \ell, m_2) \cdot \left[\frac{i}{(\ell - p)^2 - m_2^2} \right] \tau_2^\delta \left[\frac{-i \eta_{\delta\gamma}}{(\ell + q)^2} \right], \quad (\text{D3})$$

$$i\mathcal{M}_{4(e)}(q) = \tau_2^\alpha \left[\frac{-i \eta_{\alpha\beta}}{q^2} \right] \int \frac{d^4\ell}{(2\pi)^4} \tau_4^{\rho\sigma(\beta\gamma)}(q, \ell + q) \left[\frac{i\mathcal{P}_{\rho\sigma\mu\nu}}{\ell^2} \right] \tau^{\mu\nu(\delta)}(p, p', m_2) \left[\frac{-i \eta_{\delta\gamma}}{(\ell + q)^2} \right], \quad (\text{D4})$$

- Figs 5 and 6:

$$i\mathcal{M}_{5(a)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_1^{\mu\nu} \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{\ell^2} \right] \tau_3^{\alpha\beta\gamma\delta}(k, k', m_2) \left[\frac{i\mathcal{P}_{\gamma\delta\rho\sigma}}{(\ell + q)^2} \right] \tau_1^{\rho\sigma}(k - \ell, k', m_1), \quad (\text{D5})$$

$$i\mathcal{M}_{6(a)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_2^\alpha(k, \ell + k, m_1) \left[\frac{-i \eta_{\alpha\beta}}{\ell^2} \right] \tau_6^{\rho\sigma(\beta)}(p, p', m_2) \left[\frac{i\mathcal{P}_{\rho\sigma\mu\nu}}{(\ell + q)^2} \right] \tau_1^{\mu\nu}(\ell + k, k', m_1) \left[\frac{i}{(\ell + k)^2 - m_1^2} \right], \quad (\text{D6})$$

- Figs 8(a) and (c):

$$i\mathcal{M}_{8(a)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_2^\alpha(k, k + \ell, e_1) \left[\frac{-i \eta_{\alpha\beta}}{\ell^2} \right] \tau_2^\beta(p, p - \ell, e_2) \left[\frac{i}{(\ell - p)^2 - m_2^2} \right] \tau_1^{\rho\sigma}(p - \ell, p', m_2) \cdot \left[\frac{i\mathcal{P}_{\rho\sigma\mu\nu}}{(\ell + q)^2} \right] \tau_1^{\mu\nu}(\ell + k, k', m_1) \left[\frac{i}{(\ell + k)^2 - m_1^2} \right] \quad (\text{D7})$$

$$i\mathcal{M}_{8(c)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_2^\alpha(k, \ell + k, e_1) \left[\frac{-i \eta_{\alpha\beta}}{\ell^2} \right] \tau_2^\beta(p' + \ell, p', e_2) \left[\frac{i}{(\ell + p')^2 - m_2^2} \right] \tau_1^{\rho\sigma}(p, p' + \ell, m_2) \cdot \left[\frac{i\mathcal{P}_{\rho\sigma\mu\nu}}{(\ell + q)^2} \right] \tau_1^{\mu\nu}(\ell + k, k', m_1) \left[\frac{i}{(\ell + k)^2 - m_1^2} \right], \quad (\text{D8})$$

- Figs 9(a) and (b):

$$i\mathcal{M}_{9(a)}(q) = \frac{1}{2!} \int \frac{d^4\ell}{(2\pi)^4} \tau_3^{\mu\nu\alpha\beta}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\rho\sigma}}{\ell^2} \right] \tau_3^{\rho\sigma\gamma\delta}(p, p', m_2) \left[\frac{i\mathcal{P}_{\gamma\delta\alpha\beta}}{(\ell + q)^2} \right], \quad (\text{D9})$$

$$i\mathcal{M}_{9(b)}(q) = \int \frac{d^4\ell}{(2\pi)^4} \tau_6^{\mu\nu(\alpha)}(k, k', e_1) \left[\frac{-i\eta_{\alpha\beta}}{\ell^2} \right] \tau_6^{\rho\sigma(\beta)}(p, p', e_2) \left[\frac{i\mathcal{P}_{\rho\sigma\mu\nu}}{(\ell + q)^2} \right]. \quad (\text{D10})$$

- Figs 10(c), (d) and 11:

$$i\mathcal{M}_{10(c)}(q) = \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{q^2} \right] \bar{\Pi}^{\alpha\beta, \gamma\delta} \left[\frac{i\mathcal{P}_{\gamma\delta\rho\sigma}}{q^2} \right] \tau_1^{\rho\sigma}(p, p', m_2), \quad (\text{D11})$$

$$i\mathcal{M}_{10(d)}(q) = \tau_1^{\mu\nu}(k, k', m_1) \left[\frac{i\mathcal{P}_{\mu\nu\alpha\beta}}{q^2} \right] \hat{\Pi}^{\alpha\beta, \gamma\delta}(q) \left[\frac{i\mathcal{P}_{\gamma\delta\rho\sigma}}{q} \right] \tau_1^{\rho\sigma}(p, p', m_2), \quad (\text{D12})$$

$$i\mathcal{M}_{11}(q) = \tau_1^\delta(k, k', e_1) \left[\frac{-i \eta_{\delta\gamma}}{q^2} \right] \Pi^{\gamma, \tau} \left[\frac{-i \eta_{\tau\lambda}}{q^2} \right] \tau_1^\lambda(p, p', e_2) \quad (\text{D13})$$

- [1] G. 't Hooft and M. Veltman. One-loop divergencies in the theory of gravitation. *Ann. Inst. Henri Poincaré, Sec. A*, 20(1):69–94, 1974.
- [2] M. Veltman. Gravitation. In *Methods in field theory*, pages 265–327. edited by R. Balian and J. Zinn-Justin, eds., Les Houches, Session XXVIII, 1975, North Holland Publishing Company, 1976.

- [3] M. H. Goroff and A. Sagnotti. The ultraviolet behavior of einstein gravity. *Nuc. Phys. B*, 266(3-4):709–736, 1986.
- [4] S. Deser and P. van Nieuwenhuizen. Nonrenormalizability of the Quantized Einstein-Maxwell System. *Phys. Rev. Lett.*, 32(5):245–247, 1974.
- [5] S. Deser and P. van Nieuwenhuizen. One-loop divergences of quantized Einstein-Maxwell fields. *Phys. Rev.*

- D*, 10(2):401–410, 1974.
- [6] S. Deser and P. van Nieuwenhuizen. Nonrenormalizability of the quantized Dirac-Einstein system. *Phys. Rev. D*, 10(2):411–420, 1974.
 - [7] J. F. Donoghue. General relativity as an effective field theory: The leading quantum corrections. *Phys. Rev. D*, 50(6):3874–3888, 1994. [[arXiv:gr-qc/9305057v1](#)].
 - [8] J. F. Donoghue. Leading Quantum Correction to the Newtonian Potential. *Phys. Rev. Lett.*, 72(19):2996–2999, 1994. [[arXiv:gr-qc/9310024v2](#)].
 - [9] J. F. Donoghue. Introduction to the Effective Field Theory Description of Gravity. [[arxiv:gr-qc/9512024](#)], 1995. Talk given at Advanced School on Effective Theories, Almunecar, Spain, 25 Jun – 1 Jul 1995.
 - [10] H. W. Hamber and S. Liu. On the quantum corrections to the newtonian potential. *Phys. Lett. B*, 357:51–56, 1995. [[arXiv:hep-th/9505182](#)].
 - [11] A. Akhundov and A. Shiekh. A Review of Leading Quantum Gravitational Corrections to Newtonian Gravity. [[arXiv:gr-qc/0611091v2](#)], pages 1–22, 2006.
 - [12] A. Akhundov et al. Gravitational interaction to one loop in effective quantum gravity. *Phys. Lett. B*, 395:16–23, 1997. [[arXiv:hep-qc/9611018](#)].
 - [13] N. E. J. Bjerrum-Bohr et al. Quantum gravitational corrections to the nonrelativistic scattering potential of two masses. *Phys. Rev. D*, 67(084033):1–12, 2003. [[arXiv:hep-th/0211072](#)], *Erratum-ibid.* D71, 069903 (2005).
 - [14] N. E. J. Bjerrum-Bohr. Leading quantum gravitational corrections to scalar QED. *Phys. Rev. D*, 66(084023):1–11, 2002. [[arXiv:hep-th/0206236](#)].
 - [15] N. E. J. Bjerrum-Bohr et al. Quantum Corrections to the Schwarzschild and Kerr metrics. *Phys. Rev. D*, 68(084005):1–16, 2003. [[arXiv:hep-th/0211071](#)], *Erratum-ibid.* D71, 069904 (2005).
 - [16] M. S. Butt. Leading quantum gravitational corrections to QED. *Phys. Rev. D*, 74(125007):1–12, 2006. [[arxiv:hep-qc/0605137](#)].
 - [17] A. Einstein. Die Grundlage der allgemeinen Relativitätstheorie. *Ann. d. Physik*, 49(7):769–822, 1916.
 - [18] Bruce S. DeWitt. Quantum Theory of Gravity I. *Phys. Rev.*, 160(5):1113–1148, 1967. see also: Quantum Theory of Gravity II. *Phys. Rev.*, 162(5):1195–1239, 1967 and Quantum Theory of Gravity III. *Phys. Rev.*, 162(5):1239–1256.
 - [19] L. D. Faddeev and V. N. Popov. Feynman Diagrams for the Yang-Mills Field. *Phys. Lett. B*, 25(1):29–30, 1967.
 - [20] J. Schwinger. Quantized gravitational field. *Phys. Rev.*, 130(3):1253–1258, 1963. see also: Quantized Gravitational Field. II. *Phys. Rev.* 132(3):1317–1321, 1963.
 - [21] S. Weinberg. Photons and Gravitations in S-Matrix Theory: Derivation of Charge Conservation and Equality of Gravitational and Inertial Mass. *Phys. Rev.*, 135(4 B):B1049 – B1056, 1964.
 - [22] S. Mandelstam. Feynman rules for the gravitational field from the coordinate-independent field-theoretic formalism. *Phys. Rev.*, 175(5):1604–1623, 1968.
 - [23] A. E. M. van de Ven. Two-loop quantum gravity. *Nuc. Phys.*, B378:309–366, 1992.
 - [24] I. J. Muzinich and S. Vokos. Long range forces of quantum gravity. *Phys. Rev. D*, 52(6):3472–3483, 1995. [[arxiv:hep-th/9501083](#)].
 - [25] K. A. Kazakov. Notion of Potential in Quantum Gravity. *Phys. Rev. D*, 63(4):044004, 2001. [[arxiv:hep-th/0009220](#)].
 - [26] D. A. R. Dalvit et al. Running coupling constants, Newtonian potential, and nonlocalities in the effective action. *Phys. Rev. D*, 50(2):1001–1009, 1994. [[arxiv:gr-qc/9402003](#)].
 - [27] N. E. J. Bjerrum-Bohr. *Cand. Scient. thesis*. University of Copenhagen, 2001.
 - [28] D. M. Capper et al. Photon corrections to the graviton propagator. *Phys. Rev. D*, 10(2):461–467, 1974. ebd. Capper, D. M. und M. J. Duff: The One-Loop Neutrino Contribution to the Graviton Propagator. *Nuc. Phys. B*, 82(1):147–154, 1974.
 - [29] B. R. Holstein and A. Ross. Long Distance Effects in Mixed Electromagnetic-Gravitational Scattering. [arXiv:0802.0717 \[hep-ph\]](#).
 - [30] S. Faller. [arXiv:0708.1701v2 \[hep-th\]](#).
 - [31] Mathematica 5.2 is registered trademark of Wolfram Research, Inc.
 - [32] Ricci version 1.51 copyright 1992 – 2004 John M. Lee